Amendments to the Specification:

DM5109

Please replace the paragraph beginning on page 1, line 16 with the following paragraph:

In addition, it is reported that 4-aminobenzopyran derivatives that have \(\beta 3\)-receptor stimulating action and are supposed to be effective for the treatment of obesity corpulence (for example, WO 03/014113), but there has not been any mention as to the treatment of arrhythmia based on the prolongation effect on the refractory period this document.

0115101

Please replace the paragraph beginning on page 5, line 20 with the following paragraph:

- C₂₋₉ hetecyclyl group (wherein the heterocyclyl group may be arbitrarily substituted with halogen atom, C₁₋₆ alkyl group (wherein the alkyl group may be arbitrarily substituted with halogen atom, C₁₋₆ alkoxy group (wherein the alkoxy group may be arbitrarily substituted with halogen atom), amino group, carboxy group or hydroxy group), C₁₋₆ alkoxy group (wherein the alkoxy group may be arbitrarily substituted with halogen atom), C₆₋₁₄ aryl group, C₂₋₉ heteroaryl group (wherein each of the aryl group or heteroaryl group may be arbitrarily substituted with 1 to 3 R¹⁸ wherein R¹⁸ has the same above mentioned meaning as R¹⁰), hydroxy group, nitro group, cyano group, formyl group, formamide group, amino group, C₁₋₆ alkylamino group, di-C₁₋₆ alkylamino group, C₁₋₆ alkylamino group, di-C₁₋₆ alkylamino group, C₁₋₆ alkylaminocarbonyl group, di-C₁₋₆ alkylaminocarbonyl group, C₁₋₆ alkoxycarbonyl group; aminosulfonyl group, C₁₋₆ alkylsulfonyl gr

 C_{1-6} alkylamino group, di- C_{1-6} alkylamino group, C_{1-6} alkylamino group, C_{1-6} alkylaminocarbonyl group, di- C_{1-6} alkylaminocarbonyl group, C_{1-6} alkylaminocarbonyl group, C_{1-6} alkylaminocarbonyl group, C_{1-6} alkylaminocarbonyl group, C_{1-6} alkylaminosulfonyl group, C_{1-6} a

Jul 3/04

Please replace the paragraph beginning on page 9, line with the following paragraph:

(8) The benzopyran derivative or pharmaceutically acceptable salt thereof as set forth in (7), wherein R^6 is C_{6-14} aryl group wherein the aryl group may be arbitrarily substituted with 1 to 3 halogen atom or amino group, when and when a plurality of substituents are present, they may be identical or different from each other;

Dr. 104

Please replace the paragraph beginning on page 9, line 19 with the following paragraph:

forth in (5), wherein m is an integer of 1 to 3, n is 0, and R⁶ is C₂₋₄ alkyl group (wherein the alkyl group may be arbitrarily substituted with halogen atom, C₁₋₆ alkoxy group (wherein the alkoxy group may be arbitrarily substituted with halogen atom), amino group, carboxy group or hydroxy group), C₁₋₆ alkoxy group (wherein the alkoxy group may be arbitrarily substituted with halogen atom), amino group, carboxy group or hydroxy group), C₃₋₈ cycloalkyl group, C₃₋₈ cycloalkenyl group (wherein the cycloalkyl group or cycloalkenyl group may be arbitrarily substituted with halogen atom, C₁₋₆ alkyl group (wherein the alkyl group may be arbitrarily substituted with halogen atom, C₁₋₆ alkoxy group (wherein the alkoxy group may be arbitrarily substituted with halogen atom), amino group, carboxy group or hydroxy group), C₁₋₆ alkoxy group (wherein the alkoxy group may be arbitrarily substituted with halogen atom), amino group, carboxy group or hydroxy group),

group, C₁₋₆ alkoxy group (wherein the alkoxy group may be arbitrarily substituted with halogen atom) or hydroxy group), C₁₋₆ alkylcarbonyl group, aminocarbonyl group, amino group, carboxy group or cyano group;

1015109

Please replace the paragraph beginning on page 19, line 17 with the following paragraph:

7-chloro-4-[{{[2-tetrahydropyran-4-yl)ethyl}]amino}}-2,2,9-trimethyl-3,4-dihydro-2H-pyrano[2,3-g]quinolin-3-ol,

Please replace the paragraph bridging pages 23 and 24 with the following paragraph:

Examples of C₃₋₈ cycloalkyl group are such as c-propyl, c-butyl, i-methyl-c-propyl, 2-methyl-c-propyl, c-pentyl, 1-methyl-c-butyl, 2-methyl-c-butyl, 3-methyl-c-butyl, 1,2-dimethyl-c-propyl, 2,3-dimethyl-c-propyl, 1-ethyl-c-propyl, 2-ethyl-c-propyl, c-hexyl, c-heptyl, c-octyl, 1-methyl-c-hexyl, 2-methyl-c-hexyl, 3-methyl-c-hexyl, 1,2-dimethyl-c-hexyl, 2,3-dimethyl-c-propyl, 1-methyl-c-pentyl, 2-methyl-c-pentyl, 3-methyl-c-pentyl, 1-ethyl-c-butyl, 2-ethyl-c-butyl, 3-ethyl-c-butyl, 1,2-dimethyl-c-butyl, 2,3-dimethyl-c-butyl, 2,4-dimethyl-c-butyl, 3,3-dimethyl-c-butyl, 1-n-propyl-c-propyl, 2-n-propyl-c-propyl, 1-i-propyl-c-propyl, 2-i-propyl-c-propyl, 1,2,2-trimethyl-c-propyl, 1,2,3-trimethyl-c-propyl, 2,2,3-trimethyl-c-propyl, 1-ethyl-2-methyl-c-propyl, 2-ethyl-1-methyl-c-propyl, 2-ethyl-2-methyl-c-propyl, 2-ethyl-3-methyl-c-propyl, and the like.

9m 109

Please replace the paragraph beginning on page 24, line with the following paragraph:

Examples of C_{3.8} cycloalkenyl group are such as 1-c-pentenyl, 2-c-pentenyl, 3-c-pentenyl, 1-methyl-2-c-pentenyl, 1-methyl-3-c-pentenyl, 2-methyl-1-c-pentenyl, 2-methyl-2-c-pentenyl, 2-methyl-4-c-pentenyl, 2-methyl-1-c-pentenyl, 2-methyl-5-c-pentenyl, 3-methyl-1-c-pentenyl, 3-methyl-2-c-pentenyl, 3-methyl-4-c-pentenyl, 3-methyl-4-c-pentenyl, 3-methyl-5-c-pentenyl, 3-methyl-4-c-pentenyl, 1-c-hexenyl, 2-c-hexenyl, 3-c-hexenyl, 1-c-heptynyl, 2-c-heptynyl, 3-c-heptynyl, 4-c-heptynyl, 1-c-octynyl, 2-c-octenynyl, 4-c-octenynyl, 3-c-octenynyl, 3-d-octenynyl, 3-

Please replace the paragraph starting on page 29, line 1 with the following paragraph:

Examples of di-C₁₋₆ alkylaminocarbonyl group are such as dimethylaminocarbonyl, diethylaminocarbonyl, di-n-propylaminocarbonyl, di-i-propylaminocarbonyl, di-c-propylaminocarbonyl, di-n-butylaminocarbonyl, di-i-butylaminocarbonyl, di-s-butylaminocarbonyl, di-t-butylaminocarbonyl, di-1-pentylaminocarbonyl, di-2-pentylaminocarbonyl, di-3-pentylaminocarbonyl, di-i-pentylaminocarbonyl, di-c-pentylaminocarbonyl, di-c-pentylaminocarbonyl, di-1-hexylaminocarbonyl, di-2-hexylaminocarbonyl, di-3-hexylaminocarbonyl, di-(1-methyl-n-pentyl)aminocarbonyl, di-(1,1,2-trimethyl-n-propyl)aminocarbonyl, di-(1,2,2-trimethyl-n-propyl)aminocarbonyl, di-(3,3-dimethyl-n-butyl)aminocarbonyl, methyl(ethyl)aminocarbonyl, methyl(n-propyl)aminocarbonyl, methyl(i-propyl)aminocarbonyl, methyl(c-propyl)aminocarbonyl, methyl(i-butyl)aminocarbonyl, methyl(i-butyl)aminocarbonyl,

methyl(s-butyl)aminocarbonyl, methyl(t-butyl)aminocarbonyl, methyl(c-butyl)aminocarbonyl, ethyl(n-propyl)aminocarbonyl, ethyl(i-propyl)aminocarbonyl, ethyl(c-propyl)aminocarbonyl, ethyl(n-butyl)aminocarbonyl, ethyl(i-butyl)aminocarbonyl, ethyl(s-butyl)aminocarbonyl, ethyl(t-butyl)aminocarbonyl, ethyl(c-butyl)aminocarbonyl, n-propyl(i-propyl)aminocarbonyl, n-propyl(c-propyl)aminocarbonyl, n-propyl(n-butyl)aminocarbonyl, n-propyl(i-butyl)aminocarbonyl, npropyl(s-butyl)aminocarbonyl, n-propyl(t-butyl)aminocarbonyl, n-propyl(c-butyl)aminocarbonyl, i-propyl(c-butyl)aminocarbonyl i-propyl(c-<u>propyl</u>)aminocarbonyl, i-propyl(n-butyl)aminocarbonyl, i-propyl(i-butyl)aminocarbonyl, i-propyl(s-butyl)aminocarbonyl, i-propyl(t-butyl)aminocarbonyl, i-propyl(c-butyl)aminocarbonyl, c-propyl(n-butyl)aminocarbonyl. c-propyl(i-butyl)aminocarbonyl, c-propyl(s-butyl)aminocarbonyl, c-propyl(t-butyl)aminocarbonyl, c-propyl(c-butyl)aminocarbonyl, n-butyl(ibutyl)aminocarbonyl, n-butyl(s-butyl)aminocarbonyl, n-butyl(t-butyl)aminocarbonyl, n-butyl(c-butyl)aminocarbonyl, i-butyl(s-butyl)aminocarbonyl, i-butyl(t-butyl)aminocarbonyl, i-butyl(t-butyl)aminocarbonyl, i-butyl(c-butyl)aminocarbonyl, s-butyl(t-butyl)aminocarbonyl, s-butyl(c-butyl)aminocarbonyl, t-butyl(c-butyl)aminocarbonyl, and the like.

275109

Please replace the paragraph beginning on page 30, line with the following paragraph:

Examples of C₃₋₈ cycloalkylcarbonyl group are such as c-propylcarbonyl, c-butylcarbonyl, i-methyl-c-propylcarbonyl, 2-methyl-c-propylcarbonyl, c-pentylcarbonyl, 1-methyl-c-butylcarbonyl, 2-methyl-c-butylcarbonyl, 3-methyl-c-butylcarbonyl, 1,2-dimethyl-c-propylcarbonyl,

2,3-dimethyl-c-propylcarbonyl, 1-ethyl-c-propylcarbonyl, 2-ethyl-c-propylcarbonyl, c-hexylcarbonyl, c-hexylcarbonyl, c-octylcarbonyl, 1-methyl-c-hexylcarbonyl, 2-methyl-c-hexylcarbonyl, 3-methyl-c-hexylcarbonyl, 1,2-dimethyl-c-hexylcarbonyl, 1,2-dimethyl-c-propylcarbonyl, 1-ethyl-c-propylcarbonyl, 1-ethyl-c-pentylcarbonyl, 2,3-dimethyl-c-pentylcarbonyl, 3-methyl-c-pentylcarbonyl, 1-ethyl-c-butylcarbonyl, 2-methyl-c-pentylcarbonyl, 3-methyl-c-pentylcarbonyl, 1-ethyl-c-butylcarbonyl,

2-ethyl-c-butylcarbonyl, 3-ethyl-c-butylcarbonyl, 1,2-dimethyl-c-butylcarbonyl,

- 1,3-dimethyl-c-butylcarbonyl, 2,2-dimethyl-c-butylcarbonyl,
- 2,3-dimethyl-c-butylcarbonyl, 2,4-dimethyl-c-butylcarbonyl,
- 3,3-dimethyl-c-butylcarbonyl, 1-n-propyl-c-propylcarbonyl, 2-n-propyl-c-propylcarbonyl, 1-i-propyl-c-propylcarbonyl, 2-i-propyl-c-propylcarbonyl,
- 1,2,2-trimethyl-c-propylcarbonyl, 1,2,3-trimethyl-c-propylcarbonyl,
- 2,2,3-trimethyl-c-propylcarbonyl, 1-ethyl-2-methyl-c-propylcarbonyl,
- 2-ethyl-1-methyl-c-propylcarbonyl, 2-ethyl-2-methyl-c-propylcarbonyl,
- 2-ethyl-3-methyl-c-propylcarbonyl, and the like.

Please replace the paragraph beginning on page 47, line 14 with the following paragraph:

(5) The benzopyran derivative or pharmaceutically acceptable salt thereof as set forth in (3), wherein R⁶ is alkyl group, cycloalkyl group or eycloalkynyl-cycloalkenyl ring;

Please replace the paragraph beginning on page 305, line 75 with the following paragraph:

The compound of formula (1-a) or (2-a) that is the compound of formula (I) or (II) wherein A is the group of formula (5), R⁴ is hydrogen atom and R³ is hydroxy group can be

21,700

Please replace the paragraph beginning on page 316, line with the following paragraph:

Brown amorphous product

¹H-NMR(CDCl₃) δ ; 1.51(s, 6H), 2.59(d, J = 0.6 Hz, 3H), 5.90(d, J = 9.9 Hz, 1H), $\frac{6,59(d, J = 9.9 \text{ Hz}, 1\text{H})}{9.9 \text{ Hz}, 1\text{H}}$, 6.59(d, J = 9.9 Hz, 1H), 7.11(d, J = 3.6 Hz, 1H), 7.25(s, 1H), 7.68(s, 1H), 8.57(d, J = 4.4 Hz, 1H)

MS(ESI⁺)m/z; 226 [M+1]⁺

Please replace the paragraph beginning on page 318, line 6 with the following paragraph:

(Yield: 59%)

Black brown oily product

¹H-NMR(CDCl₃) δ : 1.49(s, 6H), 2.54(s, 3H), 2.62(s, 3H), 5.86(d, J = 9.9 Hz, 1H), $\frac{6,55(d, J)}{6,55(d, J)}$ = 9.9 Hz, 1H) $\frac{6.55(d, J)}{6.55(d, J)}$ = 9.9 Hz, 1H), 7.00(s, 1H), 7.20(s, 1H), 7.60(s, 1H) MS(ESI⁺)m/z; 240[M+1]⁺

Please replace the paragraph beginning on page 319, line yl with the following paragraph:

(Yield: 50%)

¹H-NMR(CDCl₃) δ; 1.50(s, 6H), 2.50(s, 3H), 2.66(s, 3H), 5.87(d, J = 9.9 Hz, 1H), 6,57(d, J = 9.9 Hz, 1H), 6.57(d, J = 9.9 Hz, 1H), 7.26(s, 1H), 7.63(s, 1H), 8.48(s, 1H) MS(ESI⁺)m/z; 240 [M+1] ⁺ PISIA

Please replace the paragraph beginning on page 326, line with the following paragraph:

Pale yellow solid

mp: 65-67°C

¹H-NMR(CDCl₃) δ ; 1.86(s, 6H), 2.70(s, 1H), 7.69-7.71(2H), 7.80(s, 1H), 8.33(d, J = 8.3 Hz, 1H), 8.45(d, J = 8.3 Hz 1H), 9.01(br s, 1H)

MS(GC)m/z MS(EI)m/z; 211 [M] +

Please replace the paragraph beginning on page 326, line 20 with the following paragraph:

Green crystal

mp: 104-107°C

¹H-NMR(CDCl₃) δ; 1.54(s, 6H), 5.89(d, J = 10.2 Hz, 1H), 6.93(d, J = 10.2 Hz, 1H), 7.50(d, J = 9.1 Hz, 1H), 7.73(br s, 1H), 8.31(d, J = 9.1 Hz, 1H), 8.74(d, J = 8.5 Hz, 1H), 9.03(br s, 1H) $\frac{MS(GC)m/z}{MS(EI)m/z}; 211[M]^{+}$

Please replace the paragraph beginning on page 339, line 2 with the following paragraph:

(Yield: 78%)

97.1%-99.1%ee; CHIRALCEL OJ-R acetonitril/methanol/0.01 M sodium chloride aqueous solution = 1/3/3, Retention time: 18.9 min.

Yellow amorphous product

¹H-NMR(CDCl₃) δ ; 1.28(s, 3H), 1.65(s, 3H), 2.59(d, J = 0.8 Hz, 3H), 3.60(d, J = 4.4 Hz, 1H), 4.13(d, J = 4.4 Hz, 1H), 7.19(s, 1H), 7.29(d, 1H), 8.02(s, 1H)

MS(ESI⁺)m/z; 276 [M+1] ⁺

Dr. 1011/109

Please replace the paragraph beginning on page 345, line 18 with the following paragraph:

Synthesis Example 26

(3R*,4S*) 4 {[2 (4 aminophenyl)ethyl]amino} -2,2,9 trimethyl 3,4 dihydro 2H-pyrano[2,3-g]quinolin 3 ol (3R*,4S*)-4-{[2-(4-aminophenyl)ethyl]amino} -7-chloro -2,2,9-trimethyl-3,4-dihydro-2H-pyrano[2,3-g]quinolin-3-ol

(Yield: 40%)

Colorless amorphous product

¹H-NMR (CDCl₃) δ : 1.23 (s, 3H), 1.55 (s, 3H), 1.58 (br s, 3H), 2.57 (s, 3H), 2.71 (t, J = 7.4 Hz, 2H), 2.85-3.05 (m, 2H), 3.11 (br s, 1H), 3.57 (d, J = 10.4 Hz, 1H), 3.84 (d, J = 10.4 Hz, 1H), 6.65 (d, J = 8.5 Hz, 2H), 7.01 (d, J = 8.5 Hz, 2H), 7.11 (s, 1H), 7.25 (s, 1H), 7.81 (s, 1H).

MS (ESI') m / z; 412 $[M+1]^+$ MS (ESI') m / z; 456 $[M+45]^+$

Please replace the paragraph bridging pages 346 and 347 with the following paragraph:

Synthesis Example 30

(3R*,4S*)-7-chloro-2,2,9-trimethyl-4-{[2-(1-piperidinyl)ethyl]amino}-3,4-dihydro-2H-pyrano[2,3-g]quinolin-3-ol

(Yield: 61%)

Pale yellow amorphous product

¹H-NMR (CDCl₃) δ : 1.29 (s, 3H), 1.58 (s, 3H), 1.60 (br s, 2H), 1.50-1.70 (m, 6H), 2.30-2.60 (m, 6H), 2,58 (s, 3H), 3.06 (t, J = 5.8 Hz, 2H) 2.58 (s, 3H), 3.06 (t, J = 5.8 Hz, 2H), 3.54 (d, J = 10.4 Hz, 1H), 3.80 (d, J = 10.4 Hz, 1H), 7.13 (s, 1H), 7.23 (s, 1H), 8.06 (s, 1H).

MS (ESI) m/z; 404 [M+1]⁺
MS (ESI) m/z; 448 [M+45]⁺

line 13

D15/69

Please replace the paragraph bridging pages 348-349 with the following paragraph:

Synthesis Example 35

(3R*,4S*)-7-chloro-2,2,9-trimethyl-4-[(2,2-diethoxyethyl)amino]-3,4-dihydro-2H-pyrano[2,3-diethoxyethyl)

g]quinolin-3-ol maleate 1-maleate

(Yield: 88%)

White solid

¹H-NMR (CD₃OD) δ: 1.23-1.30 (m, 9H), 1.57 (s, 3H), 2.64 (s, 3H), 3.50-3.85 (m, 4H), 4.02

(d, J = 10.2 Hz, 1H), 6.27 (s, 1H), 7.37 (s, 1H), 7.49 (s, 1H), 8.13 (s, 1H)

Free form

(3R*,4S*)-7-chloro-2,2,9-trimethyl-4-[(2,2-diethoxyethyl)amino]-3,4-dihydro-2H-pyrano[2,3-

g]quinolin-3-ol

Pale yellow amorphous product

 $MS (ESI^{+}) m / z; 410 [M+1]^{+}$

MS (ESI') m / z; 453 [M+45]⁺

Please replace the paragraph beginning on page 350 with the following paragraph:

Synthesis Example 40

(3R*,4S*) 7-chloro-2,2,9-trimethyl-4{[2-(2-pyridy)lethyl]amino}-3,4-dihydro-2H-

pyrano[2,3-g]quinolin-3-ol(3R*,4S*)-7-chloro-2,2,9-trimethyl-4{[2-(2-pyridyl)ethyl]amino}-

3,4-dihydro-2*H*-pyrano[2,3-g]quinolin-3-ol

(Yield: 83%)

Yellow amorphous product

Please replace the paragraph bridging pages 358 and 289 with the following paragraph:

Synthesis Example 55

(3R*,4S*)-7-chloro 2,2,9-trimethyl 4-[(2-tetrahydro-2H-thiopyran-4-ylethyl)amino]-3,4-dihydro-2H-pyrano[2,3-g]quinolin-3-ol-(3R*,4S*)-7-chloro-2,2,9-trimethyl-4-{[2-tetrahydro-2H-thiopyran-4-ylethyl]amino}-3,4-dihydro-2H-pyrano[2,3-g]quinolin-3-ol

Colorless amorphous product

(Yield: 63%)

¹H-NMR (CDCl₃) δ : 1.28 (s, 3H), 1.40-1.60 (m, 5H), 1.56 (s, 1H), 1.90-2.00 (m, 2H), 2.59 (s, 3H), 2.50-2.85 (m, 6H), 3.23 (s, 1H), 3.63 (d, J = 10.4 Hz, 1H), 3.87 (d, J = 10.4 Hz, 1H), 7.16 (s, 1H), 7.28 (s, 1H), 7.91 (s, 1H).

 $MS (ESI^{+}) m / z; 421 [M+1]^{+}$

 $MS (ESI) m / z; 465 [M+45]^{+}$

Please replace the paragraph beginning after page 361, line 1 with the following paragraph:

Under hydrogen stream at 1 atm, a solution of (3R*,4S*)-6-amino-3,4-dihydro-2,2-dimethyl-7-nitro-4-(2'-phenylethylamino)-2*H*-benzopyran-3-ol (10.0 g, 28.0 mmol) and 5% palladium carbon (AER type, 1 g) in ethanol (200 mL) was stirred at room temperature for 6 hours. Upon the completion of the reaction, the reaction solution was filtered through celite and concentrated to obtain the aimed product (yield: 98%).

Black amorphous product

¹H-NMR (CDCl₃) δ : 1.13 (s, 3H), 1.43 (s, 3H), $\frac{2.60-3.0-2.60-3.00}{6}$ (m, 4H), 2.5-3.5 (br 6H), 3.47 (d, J = 9.6 Hz, 1H), 3.51 (d, J = 9.6 Hz, 1H), 6.12 (s, 1H), 6.14 (s, 1H), 7.15-7.50 (m, 5H)

MS (ESI) m / z; $400[M+1]^+$, 327 (bp).

Please replace the paragraph beginning on page 362, line 18 with the following paragraph:

dihydro 2H-1-benzopyran-3-ol(3R*,4S*)-6,7-diamino-4-{[2-(2-fluorophenyl)ethyl]amino}-2,2-dimethyl-3,4-dihydro-2H-1-benzopyran-3-ol

(Yield: 87%)

Black amorphous product

 $MS (ESI^{+}) m / z; 346 [M+1]^{+}$

 $MS (ESI) m / z; 380 [M+45]^{+}$

Please replace the paragraph beginning on page 364, line with the following paragraph:

(8R*.9S*) {[2 (4 fluorophenyl)ethyl]amino} -7,7 dimethyl -8,9 dihydro 7H pyrano[2,3-g]quinoxalin -8 ol hydrochloride(3R*,4S*)-{[2-(4-fluorophenyl)ethyl]amino} -2,2-dimethyl-3,4-dihydro-2H-pyrano[2,3-g]quinoxalin-3-ol hydrochloride

(Yield: 95%)

Brown crystal

mp: 191-197°C (decomposition)

Please replace the paragraph beginning on page 365, line 15 with the following paragraph:

(3R*,4S*)-4-[(2-hydroxy-2-phenylethyl)amino]-2,2-dimethyl-3,4-dihydro-2H-pyrano[2,3-g]quinoxalin-3-ol

(Yield: 66%)

Two diastereomers that can not be divided separated

Gray amorphous product

¹H-NMR (CDCl₃) δ : 1.30 (s, 3H), 1.58 (s, 1.5H), 1.59 (s, 1.5H), 1.70 (br s, 3H), 2.90-3.10 (m, 2H), 3.71 (d, J= 10.5Hz, 1H), 3.95-4.05 (m, 1H),7.20-7.45 (m, 6H), 8.10 (s, 0.5H), 8.12 (s, 0.5H), 8.64 (d, J= 1.9 Hz, 1H), 8.73 (d, J= 1.9 Hz, 1H).

 $MS (ESI^{+}) m / z; 366 [M+1]^{+}$

 $MS (ESI) m / z; 410 [M+45]^{+}$

Please replace the paragraph beginning on page 367, line s with the following paragraph:

Synthesis Example 64

(3R*4S*) 2,2,7,8-tetramethyl 4 [(2-phenylethyl)amino] 3,4 dihydro 2H-pyrano[2,3-g]quinoxalin-3-ol-maleate (3R*4S*)-2,2,7,8-tetramethyl-4-[(2-phenylethyl)amino]-3,4-dihydro-2H-pyrano[2,3-g]quinoxalin-3-ol 1-maleate

Synthesis Example 64 was carried out similarly to the process of Synthesis Example 59.

Please replace the paragraph beginning after page 373, line 19 with the following paragraph:

6-methoxy-2,2-dimethyl-2H-1-benzopyran

A solution of 4-(1,1-dimethyl-2-propenyloxy)anisole 4-(1,1-dimethyl-2-propenyloxy)anisole in 1,2-dichlorobenzene (50 mL) was stirred at 190°C for 2 hours. Upon the completion of the reaction, the solvent was distilled off under a reduced pressure. The residue was purified by column chromatography (hexane/chloroform = 3/1) and the aimed product was obtained as red oily substance (2-step, yield: 61%).

¹H-NMR (CDCl₃) δ: 1.41 (s, 6H), 3.75 (s, 3H), 5.64 (d, *J*=9.9 Hz, 1H), 6.28 (d, *J*=9.9 Hz, 1H), 6.55 (d, *J*=2.7 Hz, 1H), 6.64-6.73 (m, 2H)

 $LC/MS (ESI^{+})m/z$: 191[M⁺+1]

Please replace the paragraph beginning after page 374, line 2 with the following paragraph:

A mixed solution of acetic acid (6.2 mL) and acetic anhydride (6.2 mL) containing 6-methoxy-2,2-dimethyl-2H-1-benzopyran (3.1 g, 16.4 mmol) was cooled with ice, nitric acid (1.37 mL, 18.0 mmol) was added dropwise and then the mixture was stirred at 0°C for 1 hour. Upon the completion of the reaction, an aqueous solution of 1 mol/L sodium hydroxide was added to the reaction solution, the resulting solution was extracted with ethyl acetate (150 mL). The organic phase was washed twice with 1 mol/L sodium hydroxide aqueous solution and once with saturated sodium chloride solution. Then, the organic phase was dried over anhydrous magnesium sulfate. After distilling off the solvent, the residue was purified by

column chromatography (hexane/ethyl acetate = 6/1) and the aimed product was obtained as yellow crystal (yield: 79%).

¹H-NMR (CDCl₃) δ: 1.44 (s, 6H), 3.91 (s, 3H), 5.85 (d, *J*=9.6 Hz, 1H), 6.33 (d, *J*=9.6 Hz, 1H), 6.69 (s, 1H), 7.34 (s, 1H)

LC/MS (ESI+): 236 [M++1] LC/MS (ESI+)m/z: 236 [M++1]

Please replace the paragraph bridging pages 374 and 375 with the following paragraph:

¹H-NMR (CDCl₃) δ: 1.26 (s, 3H), 1.58 (s, 3H), 3.53 (d, *J*=4.3 Hz, 1H), 3.90 (d, *J*=4.3 Hz, 1H), 3.95 (s, 3H), 7.08 (s, 1H), 7.33 (s, 1H)

MS (EI): 251 [M⁺] MS (EI)m/z: 251 [M⁺]

HPLC: 18.6 min (enantiomer 24.1 min)

HPLC condition: chiralcel OJ-RH, MeCN/MeOH/0.01 M NaCl aq. = 1/3/5, 1.0 ml/min, 40°C, 256 nm

Please replace the paragraph beginning on page 375, line with the following paragraph:

To a solution of (3R*, 4S*)-3,4-epoxy-6-methoxy-2, 2-dimethyl-7-nitro-3,4-dihydro-2H-1-benzopyran (2.50 g, 9.95 mmol) in 1,4-dioxane (5.0 mL), lithium perchlorate (1.06 g, 9.95 mmol) and 4-(phenylethyl)amine (1.50 mL, 11.9 mmol) were added at room temperature and the mixture was stirred at 80 °C for 1 hour. Upon the completion of the reaction, an aqueous solution of saturated ammonium chloride was added to the reaction solution, and the resulting solution was extracted with ethyl acetate. The organic phase was washed with saturated sodium chloride solution, and then dried over anhydrous sodium sulfate. After distilling off the solvent, the residue was purified by column chromatography (hexane/ethyl

and the resulting mixture was stirred at room temperature for 10 minutes. Upon the completion of the reaction, the resulting crystal was filtered off and the aimed product was obtained (yield: 93%).

15/15/19

Please replace the paragraph beginning on page 382, line with the following paragraph:

Synthesis Example 74

7-Hydroxy-6,6-dimethyl-8-(2-phenylethylamino)-7,8-dihydro-1H,6H-4,5-dioxa-1-aza-anthracene-2-one

2-methoxymethoxy-4 (1,1-dimethyl 2-propionyloxy)-1-nitro-benzene

(1,1-dimethyl-2-propynyloxy)-1-nitro-benzene

Please replace the paragraph beginning on page 383, line 3 with the following paragraph:

A solution of 2 methoxymethoxy 4 (1,1 dimethyl-2 propionyloxy) 1 nitro benzene 2-methoxymethoxy-4-(1,1-dimethyl-2-propynyloxy)-1-nitro-benzene (2.1 g, 7.92 mmol) in dichlorobenzene 1,2-dichlorobenzene (21 mL) was stirred at 20°C for 0.5 hour. Upon the completion of the reaction, the resulting mixture was concentrated and purified by silica gel column (hexane/ethyl acetate = 5/1). Thereby, a mixture (1:1) of the aimed product and the

¹H-NMR (CDCl₃) δ : 1.17 (s, 3H), 1.45 (s, 3H), 2.75-3.00 (m, 4H), 3.43 (d, J = 9.9 Hz, 1H), 3.50 (s, 3H), 3.59 (d, J = 9.9 Hz, 1H), 4.20 (s, 2H), 5.19 (s, 2H), 6.61 (s, 1H), 7.15-7.30 (m, 5H), 8.14 (s, 1H), 8.73 (s, 1H).

 $MS (ESI^{+}) m / z: 449 [M+1]^{+}$

 $MS (ESI) m / z: 447 [M-1]^{+}$

Please replace the paragraph beginning on page 386, line 1/2 with the following paragraph:

2-Chloro N-[(±) trans-3,7 dihydroxy-2,2 dimethyl 4-[(2 phenylethyl)amino] 3,4-dihydro-2H-1-benzopyran-6-yl]-acetamide 2-chloro-N-{(±)-trans-3,7-dihydroxy-2,2-dimethyl-4-[(2-phenylethyl)amino]-3,4-dihydro-2H-1-benzopyran-6-yl}-acetamide

Please replace the paragraph beginning on page 386, line 16 with the following paragraph:

To a solution of 2-chloro-N-[(±) trans 3-hydroxy 7-methoxymethoxy 2,2 dimethyl 4[(2-phenylethyl)amino] 3,4-dihydro-2H 1-benzopyran-6-yl] acetamide-2-chloro-N-{(±)trans-3-hydroxy-7-methoxymethoxy-2,2-dimethyl-4-[(2-phenylethyl)amino]-3,4-dihydro-2H1-benzopyran-6-yl}-acetamide (228 mg, 0.51 mmol) in methylene chloride (6 mL), boron
tribromide (1 M solution in methylene chloride, 2.42 mL, 2.42 mmol) was added at 0°C, and
the resulting mixture was stirred for 2 hours. Upon the completion of the reaction, methanol
and saturated sodium hydrogencarbonate aqueous solution were added thereto, and the
resulting solution was extracted with ethyl acetate, washed with saturated sodium
hydrogencarbonate aqueous solution and then with saturated sodium chloride solution, dried
over magnesium sulfate and concentrated to obtain the aimed product (yield: 100%).
Colorless amorphous product

الارادام

Please replace the paragraph beginning on page 391, line 1/6 with the following paragraph:

This compound was synthesized according to the process of Synthesis Example 19. (Yield: 30%)

Orange amorphous product

¹H-NMR (CDCl₃) δ: 1.19 (s, 3H), 1.50 (s, 3H), 2.05-2.15 (br, 2H), 2.49 (s, 3H), 3.09-3.32 (m, 10H), 4.60-5.20 (br, 2H), 7.06 (s, 1H), 7.11 (s, 1H), 7.88 (s, 1H)

MS (ΕΓ⁺) (ESI⁺) m / z; 390[M+1]⁺

Please replace the paragraph bridging pages 393 and 394 with the following paragraph:

To a solution of (3R*,4R*)-3-hydroxy-2,2,9-trimethyl-4-[(2-phenylethyl)amino]-3,4-dihydro-2H-pyrano[2,3-g]quinoline-7-carbonitrile described in Synthesis Example 14 (465 mg, 1.20 mmol) in ethanol (5 mL), sodium hydroxide aqueous solution (3 mol/L, 5 mL) was added at room temperature, and the resulting mixture was stirred for 2 hours with reflux under heating. After cooling to room temperature, the resulting solution was neutralized with 1 mol/L hydrochloric acid, precipitated brown solid was filtered off and the aimed product was obtained (yield: 90%).

Brown solid

¹H-NMR (CDCl₃) δ: 1.07 (s, 3H), 1.41 (s, 3H), 2.46 (s, 3H), 2.89-3.08 (br, 2H), 3.10-3.28 (br, 2H), 4.03-4.22 (br, 1H), 4.30-4.44 (br, 1H), 7.01-7.54 (m, 7H), 7.86 (s, 1H), 8.51-8.73 (br, 1H)

 $MS (EI^{+}) (ESI^{+}) m / z; 407 [M+1]^{+}$

3/15/67

Please replace the paragraph beginning after page 404, line with the following paragraph:

_(3R*,4S*)-7-chloro-2,2,9-trimethyl-6λ5-oxy-4-pentylamino-3,4-dihydro-2*H*-pyrano[2,3-g]quinolin-3-ol hydrochloride (3R*,4S*)-7-Chloro-2,2,9-trimethyl-6λ5-oxy-4-pentylamino-3,4-dihydro-2*H*-pyrano[2,3-g]quinolin-3-ol hydrochloride (yield: 60%).

Pale yellow crystal

mp;226-230°C (decomposition)

¹H-NMR(DMSO-d₆) δ; 0.86(t, J = 6.3 Hz, 3H), 1.16(s, 3H), 1.27-1.29(m, 4H), 1.50(s, 3H), 1.60-1.72(m, 2H), 2.54(s, 3H), 2.86(brs, 1H), 3.07(brs, 1H), 4.07-4.10(m, 1H), 4.71(d, J = 8.5Hz, 1H), 6.51(d, J = 4.7 Hz, 1H), 7.47(s, 1H), 7.67(s, 1H), 9.04(s, 1H), 9.19(brs, 1H), 9.74(brs, 1H)

 $MS(ESI^{+})m/z$; 379, 381 $[M+1]^{+}$

MS(ESI')m/z; 423, 425 [M+45]⁺

Please replace the paragraph beginning on page 405, line 1 with the following paragraph:

To a solution of t-butyl (2-phenylethyl) (3R*,4S*)-7-amino-3-hydroxy-6-methoxy-2,2-dimethyl-3,4-dihydro-2H-1-benzopyran-4-yl carbamate described in Synthesis Example 71 (1.04 g, 2.35 mmol) in pyridine (1.90 mL, 23.5 mmol), chloromethanesulfonylchloride (0.31 mL, 3.52 mmol) was added, and the resulting mixture was stirred at room temperature for 10 hours. Upon the completion of the reaction, 1 mol/L hydrochloric acid aqueous solution (ca. 30 mL) was added thereto to adjust pH to about 7, and then the resulting solution was extracted with ethyl acetate, washed with saturated sodium chloride aqueous solution, and dried over anhydrous sodium sulfate and concentrated. The resulting mixture was purified by

Amendments to the Specification:

2325

Susia

Please replace specification page 360, lines 24/26, with the following, amended lines: (3R*,4S*) 2,2 dimethyl-4-[(2 phenylethyl)amino]-3,4-dihydro-2H pyrano[2,3-g]quinoxalin-3-ol-1 maleate

(3R*,4S*)-7,7-dimethyl-9-[(2-phenylethyl)amino]-8,9-dihydro-7H-pyrano[2,3-g]quinoxalin-8-ol 1 maleate

Please replace specification page 361, lines 15-16, with the following, amended lines:

(3R*,4S*)-2,2-dimethyl-4-[(2-phenylethyl)amino]-3,4-dihydro-2H-pyrano[2,3-g]quinoxalin-3-ol (3R*,4S*)-7,7-dimethyl-9-[(2-phenylethyl)amino]-8,9-dihydro-7H-pyrano[2,3-g]quinoxalin-8-ol

Please replace specification page 362, lines 4-5, with the following, amended lines:

(3R*,4S*)-2,2 dimethyl 4 [(2-phenylethyl)amino] 3,4 dihydro 2H-pyrano[2,3g]quinoxalin-3 ol 1 maleate (3R*,4S*)-7,7-dimethyl-9-[(2-phenylethyl)amino]-8,9-dihydro7H-pyrano[2,3-g]quinoxalin-8-ol 1 maleate

Amendments to the Specification:

10/15/09

Please replace the paragraph beginning on page 349, line of the specification with the following, amended paragraph:

(3R*,4S*) 7-chloro-2,2,9-trimethyl-4-[2 (1-pyrazolylethyl)amino] 3,4-dihydro-2H-pyrano[2,3-g]quinolin-3-ol

(3R*,4S*)-4-[2-(1H-pyrazol-1-yl)ethylamino]-7-chloro-2,2,9-trimethyl-3,4-dihydro-2H-pyrano[2,3-g]quinolin-3-ol

Dr. 5109

Please replace the paragraph beginning on page 360, line $\frac{1}{2}$ 4 with the following, amended paragraph:

(3R*,4S*)-7,7-dimethyl-9-[(2-phenylethyl)amino]-8,9-dihydro-7H-pyrano[2,3-g]quinoxalin-8-ol-1-maleate

(8R*,9S*)-7,7-dimethyl-9-[(2-phenylethyl)amino]-8,9-dihydro-7H-pyrano[2,3-g]quinoxalin-8-ol 1 maleate

800/5/06

Please replace the paragraph beginning on page 361, line 1/5 with the following, amended paragraph:

(3R*,4S*) 7,7-dimethyl-9-[(2-phenylethyl)amino] 8,9-dihydro-7H-pyrano[2,3-g]quinoxalin-8-ol

(8R*,9S*)-7,7-dimethyl-9-[(2-phenylethyl)amino]-8,9-dihydro-7H-pyrano[2,3-g]quinoxalin-8-ol

10/18/04

Please replace the paragraph beginning on page 362, line 4 with the following, amended paragraph:

(3R*,4S*)-7,7-dimethyl-9-[(2-phenylethyl)amino] 8,9-dihydro 7H-pyrano[2,3-g]quinoxalin-8-ol-1-maleate

(8R*,9S*)-7,7-dimethyl-9-[(2-phenylethyl)amino]-8,9-dihydro-7H-pyrano[2,3-g]quinoxalin-8-ol 1 maleate

Dm

Please replace the paragraph beginning on page 362, line with the following, amended pargraph:

(3R*,4S*) 7,7 dimethyl 9 [(2 phenylethyl)amino] 8,9 dihydro 7H pyrano[2,3 g]quinoxalin-8 ol

(8R*,9S*)-7,7-dimethyl-9-[(2-phenylethyl)amino]-8,9-dihydro-7H-pyrano[2,3-g]quinoxalin-8-ol (1.18 g, 3.38 mmol) in ethyl acetate (22 mL), maleic acid (471 mg, 4.06 mmol) was added at room temperature, and the resulting mixture was stirred for 10 minutes. Upon the completion of the reaction, solid product was filtered off, washed with ethyl acetate and dried to obtain the aimed product (yield: 61%).

Pale gray crystal

Jul 2/01

Please replace the paragraph beginning on page 362, line with the following, amended paragraph:

(3R*,4S*) 4-{[2 (2-fluorophenyl)ethyl]amino}-2,2-dimethyl 3,4-dihydro-2H-pyrano[2,3-g]quinoxalin-3-ol-hydrochloride

(8R*,9S*)-9-{[2-(2-fluorophenyl)ethyl]amino}-7,7-dimethyl-8,9-dihydro-7H-pyrano[2,3-g]quinoxalin-8-ol hydrochloride

projou

Please replace the paragraph beginning on page 363, line with the following, amended paragraph:

(3R*,4S*) 4-{[2 (2 fluorophenyl)ethyl]amino}-2,2-dimethyl 3,4-dihydro-2H-pyrano[2,3-g]quinoxalin 3-ol

(8R*,9S*)-9-{[2-(2-fluorophenyl)ethyl]amino}-7,7-dimethyl-8,9-dihydro-7H-pyrano[2,3-g]quinoxalin-3-ol

10/13/10

Please replace the paragraph beginning on page 363, line with the following, amended paragraph:

(3R*,4S*) 4-{[2 (2-fluorophenyl)ethyl]amino}-2,2-dimethyl-3,4-dihydro-2H-pyrano[2,3-g]quinoxalin-3-ol-hydrochloride

(8R*,9S*)-9-{[2-(2-fluorophenyl)ethyl]amino}-7,7-dimethyl-8,9-dihydro-7H-pyrano[2,3-g]quinoxalin-3-ol hydrochloride

2112/an

Please replace the paragraph beginning on page 363, line 24 with the following, amended paragraph:

(3R*,4S*) 9 {[2 (4-fluorophenyl)ethyl]amino}-7,7-dimethyl-8,9-dihydro-7H-pyrano[2,3-g]quinoxalin-8-ol hydrochloride

(8R*,9S*)-9-{[2-(4-fluorophenyl)ethyl]amino}-7,7-dimethyl-8,9-dihydro-7H-pyrano[2,3-g]quinoxalin-8-ol hydrochloride

9/15/64

Please replace the paragraph beginning on page 364, line 1/2 with the following, amended paragraph:

(3R*,4S*) 9-{[2 (4-fluorophenyl)ethyl]amino} -7,7-dimethyl-8,9-dihydro-7H-pyrano[2,3-g]quinoxalin-8-ol

(8R*,9S*)-9-{[2-(4-fluorophenyl)ethyl]amino}-7,7-dimethyl-8,9-dihydro-7H-pyrano[2,3-g]quinoxalin-8-ol

60/12/04

Please replace the paragraph beginning on page 364, line 2/2 with the following, amended paragraph:

(8R*,9S*)-9-{[2-(4-fluorophenyl)ethyl]amino} 7,7-dimethyl-8,9-dihydro-7H-pyr-ano[2,3-g]quinoxalin-8-ol hydrochloride

(8R*,9S*)-9-{[2-(4-fluorophenyl)ethyl]amino}-7,7-dimethyl-8,9-dihydro-7H-pyrano[2,3-g]quinoxalin-8-ol hydrochloride

Please replace the paragraph beginning on page 365, line 2 with the following, amended paragraph:

(3R*,4S*) 4 [(2-hydroxy-2-phenylethyl)amino]-2,2-dimethyl-3,4-dihydro 2H pyrano[2,3-g]quinoxalin-3-ol

(8R*,9S*)-9-[(2-hydroxy-2-phenylethyl)amino]-7,7-dimethyl-8,9-dihydro-7H-pyrano[2,3-g]quinoxalin-8-ol

2013/04

Please replace the paragraph beginning on page 365, line with the following, amended paragraph:

(3R*,4S*) 4 [(2 hydroxy 2 phenylethyl)amino] 2,2 dimethyl 3,4 dihydro 2H pyrano[2,3 g]quinoxalin 3 ol

(8R*,9S*)-9-[(2-hydroxy-2-phenylethyl)amino]-7,7-dimethyl-8,9-dihydro-7H-pyrano[2,3-g]quinoxalin-8-ol

Please replace the paragraph beginning on page 366, line 2 with the following, amended paragraph:

(3R*,4S*) 2,2 dimethyl 4-pentylamino 3,4 dihydro-2H-pyrano[2,3-g]quinoxalin-3-ol hydrochloride

(8R*,9S*)-7,7-dimethyl-9-(pentylamino)-8,9-dihydro-7H-pyrano[2,3-g]quinoxalin-8-ol hydrochloride

2015/09

Please replace the paragraph beginning on page 366, line with the following, amended line:

(3R*,4S*) 2,2-dimethyl-4-pentylamino-3,4-dihydro-2H-pyrano[2,3-g]quinoxalin-3-ol (8R*,9S*)-7,7-dimethyl-9-(pentylamino)-8,9-dihydro-2H-pyrano[2,3-g]quinoxalin-8-ol

2015/04

Please replace the paragraph beginning on page 366, line 25 with the following, amended paragraph:

(3R*,4S*) 2,2 dimethyl 4 pentylamino 3,4 dihydro 2H-pyrano[2,3-g]quinoxalin 3-ol hydrochloride

(8R*,9S*)-7,7-dimethyl-9-(pentylamino)-8,9-dihydro-2H-pyrano[2,3-g]quinoxalin-8-ol

30/10/04

Please replace the paragraph beginning on page 367, line 8 with the following, amended paragraph:

(3R*4S*) 2,2,7,8 tetramethyl 4-[(2 phenylethyl)amino] 3,4 dihydro 2H-pyrano[2,3-g]quinoxalin-3-ol-maleate

(8R*,9S*)-2,3,7,7-tetramethyl-9-[(2-phenylethyl)amino]-8,9-dihydro-7H-pyrano[2,3-g]quinoxalin-8-ol maleate

10/15/04

Please replace the paragraph beginning on page 367, line 1/4 with the following, amended paragraph:

3R*4S*)-2,2,7,8-tetramethyl-4-[(2-phenylethyl)amino] 3,4-dihydro-2H-pyrano[2,3-g]quinoxalin-3-ol

(8R*9S*)-2,3,7,7-tetramethyl-9-[(2-phenylethyl)amino]-8,9-dihydro-7H-pyrano[2,3-g]quinoxalin-8-ol

pmiolisher

Please replace the paragraph beginning on page 367, line 26 with the following, amended paragraph:

(3R*,4S*) 7,8 diethyl 2,2 dimethyl 4 [(2-phenylethyl)amino] 3,4 dihydro-2H-pyrano[2,3-g]quinoxalin-3 ol

(8R*,9S*)-2,3-diethyl-7,7-dimethyl-9-[(2-phenylethyl)amino]-8,9-dihydro-7H-pyrano[2,3-g]quinoxalin-8-ol

2011/3/04

Please replace the paragraph beginning on page 368, line 1/2 with the following, amended paragraph:

(3R*,4S*) 2,2,8-trimethyl-7-phenyl-4-[(2-phenylethyl)amino] 3,4-dihydro 2H-pyrano[2,3-g]quinoxalin-3-ol

(8R*,9S*)-3,7,7-trimethyl-9-[(2-phenylethyl)amino]-2-phenyl-8,9-dihydro-7H-pyrano[2,3-g]quinoxalin-8-ol

10/1/109

Please replace the paragraph beginning on page 368, line 25 with the following, amended paragraph:

(3R*,4S*) 2,2,7-trimethyl-8-phenyl-4-[(2-phenylethyl)amino] 3,4-dihydro-2H-pyrano[2,3-g]quinoxalin-3-ol

(8R*,9S*)-2,7,7-trimethyl-9-[(2-phenylethyl)amino]-3-phenyl-8,9-dihydro-7H-pyrano[2,3-g]quinoxalin-8-ol

Dullon

Please replace the paragraph beginning on page 369, line in with the following, amended paragraph:

(3R*,4S*)-2,2,8 trimethyl-4-[(2-phenylethyl)amino]-3,4 dihydro-2H-pyrano[2,3-g]quinoxalin-3-ol-1 maleate

(8R*,9S*)-3,7,7-trimethyl-9-[(2-phenylethyl)amino]-8,9-dihydro-7H-pyrano[2,3-g]quinoxalin-8-ol 1 maleate

2011310

Please replace the paragraph beginning on page 369, line 16 with the following, amended paragraph:

(3R*,4S*) 2,2,8-trimethyl-4-[(2-phenylethyl)amino] 3,4-dihydro-2H-pyrano[2,3-g]quinoxalin-3-ol

(8R*,9S*)-3,7,7-trimethyl-9-[(2-phenylethyl)amino]-8,9-dihydro-7H-pyrano[2,3-g]quinoxalin-8-ol

July A

Please replace the paragraph beginning on page 370, line with the following, amended paragraph:

(3R*,4S*) 4-[(2-eyelohexylethyl)amino)]-2,2-dimethyl-3,4-dihydro-2H-pyrano[2,3-g]quinoxalin-3-ol hydrochloride

(8R*,9S*)-9-[(2-cyclohexylethyl)amino)]-7,7-dimethyl-8,9-dihydro-7H-pyrano[2,3-g]quinoxalin-8-ol hydrochloride

975/64

Please replace the paragraph beginning on page 370, line with the following, amended paragraph:

(3R*,4S*) 4-amino 2,2-dimethyl-3,4-dihydro 2*H*-pyrano[2,3-g]quinoxalin-3-ol (8R*,9S*)-9-amino-7,7-dimethyl-8,9-dihydro-7*H*-pyrano[2,3-g]quinoxalin-8-ol

holision holision

Please replace the paragraph beginning on page 370, line 26 with the following, amended paragraph:

(3R*,4S*) 4-[(2-eyelohexylethyl)amino)] 2,2-dimethyl-3,4-dihydro-2*H*-pyrano[2,3-g]quinoxalin-3-ol

(8R*,9S*)-9-[(2-cyclohexylethyl)amino)]-7,7-dimethyl-8,9-dihydro-7H-pyrano[2,3-g]quinoxalin-8-ol

Please replace the paragraph beginning on page 371, line 3 with the following, amended paragraph:

(4R*,4S*) 4 amino 2,2 dimethyl 3,4 dihydro 2H pyrano[2,3 g]quinoxalin 3 ol (8R*,9S*)-9-amino-7,7-dimethyl-8,9-dihydro-7H-pyrano[2,3-g]quinoxalin-8-ol (100 mg, 0.408 mmol) in methanol (2 mL), cyclohexylmethyl aldehyde (103 mg, 0.816 mmol) was added, and the resulting mixture was stirred at room temperature for 20 minutes. Sodium cyanoborohydride (51 mg, 0.816 mmol) was added thereto, and the resulting mixture was stirred at room temperature for 1 hour. Upon the completion of the reaction, saturated sodium hydrogenearbonate aqueous solution was added thereto, the resulting solution was extracted with ethyl acetate, washed with saturated sodium chloride solution, and dried over magnesium sulfate and concentrated. The resulting mixture was purified by silica gel column (hexane/ethyl acetate = 2/1) to obtain the aimed product (yield: 48%).

9/15/09

Please replace the paragraph beginning on page 371, line 19 with the following, amended paragraph:

(3R*,4S*) 4-{(2-cyclohexylethyl)amino}-2,2-dimethyl-3,4-dihydro-2H-pyrano{2,3-g}quinoxalin-3-ol-hydrochloride

(8R*,9S*)-9-[(2-cyclohexylethyl)amino]-7,7-dimethyl-8,9-dihydro-7H-pyrano[2,3-g]quinoxalin-8-ol hydrochloride

10/15/04

Please replace the paragraph beginning on page 371, line 28 with the following, amended paragraph:

(±) trans-3-hydroxy-2,2-dimethyl-4-[(2-phenylethyl)amino]-2,3,4,6-tetrahydro-pyrano[2,3-f]benzimidazol-7one

(±)-trans-7-hydroxy-6,6-dimethyl-8-[(2-phenylethyl)amino]-3,6,7,8-tetrahydrochromeno[7,6-d]imidazol-2(1H)-one